

Structural and dynamic studies of naphtha crude residue with different chemical nature

Kemalov A., Kemalov R.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

It is known that the tars of heavy oils of naphthene-aromatic base with high concentrations of resin-asphaltene substances (RAS) and insufficient content of high-molecular paraffins are the most suitable raw material for production of a wide range of bituminous products in general and bituminous insulating material (BIM). Naphtha crude residues (NCR) of paraffin and high-paraffin crude oils are usually used due to limited oil production. It is notable that the production of refractory special bitumens at high-temperature oxidation of residual materials results in production of non-hydrocarbon compounds - carbenes and carboides that are insoluble in hot benzene, as well as the compounds of paraffin-naphthenic base containing in extracted oil. Considering the NCR as a thermodynamic system of phases A, B and C, it should be noted that the structure of carbon links of phases can be assumed by evaluation of molecular mobility and structural-dynamic parameters. Structural-dynamic states of NCR at different temperatures were studied using relaxometer NMR 08BK/RS and used as raw material for oxidation. © IDOSI Publications, 2013.

<http://dx.doi.org/10.5829/idosi.wasj.2013.22.tt.22147>

Keywords

IR - spectroscopy, Naphtha crude residues, Nuclear magnetic resonance relaxation (NMR-relaxation), Oxidative polymerization, Paraffin-asphalt associates, Physical-chemical modification, Structurally complex links